
The Audubon Society
Field Guide to
North American Reptiles and
Amphibians

A Chanticleer Press Edition

John L. Behler, Curator of Herpetology,
New York Zoological Society, and
F. Wayne King, Director,
Florida State Museum

Alfred A. Knopf, New York

This is a Borzoi Book
Published by Alfred A. Knopf, Inc.

All rights reserved. Copyright 1979
under the International Union for the
protection of literary and artistic
works (Berne). Published in the United
States by Alfred A. Knopf, Inc., New
York, and simultaneously in Canada by
Random House of Canada Limited,
Toronto. Distributed by Random
House, Inc., New York.

Prepared and produced by
Chanticleer Press, Inc., New York.

Color reproductions by Nievergelt
Repro AG, Zurich, Switzerland.
Printed and bound by Amilcare Pizzi,
S.p.A., Milan, Italy.
Type set in Garamond by
Dix Typesetting Co. Inc., Syracuse,
New York.

First Printing

QL651.K56 1979
598.1'097 79-2217
ISBN 0-394-50824-6

Trademark "Audubon Society" used by
publisher under license from the
National Audubon Society, Inc.

John L. Behler is Curator of
Herpetology at the New York
Zoological Society's Bronx Zoo. Since
he joined the staff of the Society in
1970, his popular articles on reptiles
and amphibians have appeared regularly
in *Animal Kingdom* magazine. In
addition to his ongoing work with
native northeastern reptiles and
amphibians, Behler has participated in
numerous national and international
studies, including the United Nations
Development Programme's 1975
expedition to Papua, New Guinea, to
examine that country's crocodylian
conservation program.

F. Wayne King is Director of the
Florida State Museum and an
internationally recognized expert on the
conservation of endangered animals,
especially reptiles and amphibians. He
is Deputy Chairman for the Species
Survival Commission of the
International Union for the
Conservation of Nature and Natural
Resources, former Director of
Conservation for the New York
Zoological Society, and a past Curator
of Herpetology at the New York
Zoological Society's Bronx Zoo. He has
written numerous scientific and popular
articles on reptiles and amphibians as
well as other wildlife.

Mexican (*L. g. nigritus*), uniformly black or black with small light spot on each side scale and some back scales, 23–25 scale rows; w. Sonora and extreme nw. Sinaloa, Mexico, into se. Arizona.

Desert (*L. g. splendida*), back dark brown or black with narrow light crossbands, scales on sides have central light spot, 23–25 scale rows; c. Texas west to se. Arizona south to c. Mexico.

Active during the day, especially early in the morning or near dusk, but becomes nocturnal in the warm summer months. It is primarily terrestrial, occasionally climbing into shrubs. A strong constrictor, it eats snakes—including rattlesnakes, Copperheads, and coral snakes—as well as lizards, mice, birds, and eggs. Record longevity exceeds 24 years.

601, 602, 603 **Gray-banded Kingsnake**
(*Lampropeltis mexicana*)

Description: 24–47½" (61–120.7 cm). *Highly variable pattern* of white-bordered gray crossbands alternating with black-bordered reddish-orange crossbands or blotches. Head distinct from neck; eyes large. Scales smooth, in 25 rows. Anal plate single.

Breeding: Mates in spring. 4–9 eggs, 1¼–1¾" (32–45 mm) long, are laid late May to late July, hatch in 9–11 weeks. Young are 9–11" (23–28 cm) long.

Habitat: Arid mesquite-creosote bush desert flats, barren rocky hillsides, canyons, limestone ledges, ranging into semiarid mountainous situations; 1,200–7,500' (350–2,300 m).

Range: Trans-Pecos region east to Balcones Escarpment of Edwards Plateau of Texas, south into Mexico.

Subspecies: Three, poorly defined; 1 enters our range, *L. m. alterna*.



Once considered rare, this handsome snake became a favorite of collectors. A secretive, nocturnal species it proved to be considerably more common than believed. Lizards are its staple diet, but it also eats frogs, small snakes, and rodents.

598 **Sonora Mountain Kingsnake**
(*Lampropeltis pyromelana*)

Subspecies: Arizona, Utah, Huachuca

Description: 20–41" (50.8–104.1 cm). Tricolored kingsnake with red, black, and white, cream, or yellow bands. *Light rings do not widen near belly scales*. Black bands border red and light bands and become narrow or disappear on sides. Top of head black; *snout light colored*. Scales smooth, in 23–25 rows. Anal plate single.

Breeding: Mates in spring. Clutches of 3–6 elongated eggs, 1½–2¾" (44–60 cm) long, are deposited June to July and hatch in about 2½–3 months. Young 9–10" (8.8–10 cm) long.

Habitat: Chaparral woodland and pine forests in mountainous regions; brushy rocky canyons, talus slopes, and near streams and springs; 2,800–9,100' (850–2,800 m).

Range: Nc. Arizona south into Mexico; isolated populations in Utah, Nevada, and Arizona.

Subspecies: Four; 3 in our range.

Arizona (*L. p. pyromelana*), usually more than 43 light rings on body, 10 lower lip scales; c. and se. Arizona and sw. New Mexico into nw. Mexico.

Utah (*L. p. infralabialis*), half or more of white body rings cross belly, 9 lower lip scales; e. Nevada, c. Nevada to Grand Canyon area of nw. Arizona.

Huachuca (*L. p. woodini*), usually fewer than 43 light rings on body, 10 lower lip scales; Huachuca Mountains, s. Arizona into Mexico.





601 Gray-banded Kingsnake, 24–47½", p. 620



602 Gray-banded Kingsnake, 24–47½", p. 620



603 Gray-banded Kingsnake, 24–47½", p. 620