


# The Venomous Reptiles of Latin America



**JONATHAN A. CAMPBELL and WILLIAM W. LAMAR**

*Department of Biology, The University of Texas at Arlington*

**Comstock Publishing Associates**

A DIVISION OF CORNELL UNIVERSITY PRESS  
ITHACA AND LONDON



Fig. 484. *Geophis duellmani*, male, 192 mm TL (UTA R-12333); 6.8 km south of Vista Hermosa (north slope of the Sierra de Juárez), Oaxaca, Mexico, 1661 m elevation.



Fig. 485. *Geophis semidoliatus*, female, 314 mm TL (UTA R-19910); Parque Moctezuma, Barranca de San Miguel, Cuautlalpan, Veracruz, Mexico.



Fig. 486. *Gyalopion quadrangulare*; Carretera Estatal (State Highway) 23, Sinaloa, Mexico. Photo by R. Wayne Van Devender.



Fig. 487. *Hydrodynastes bicinctus*, male, approximately 1,500 mm TL; El Tapón, Caño La Hormiga, Vichada, Colombia.



Fig. 488. *Hydrops triangularis venezuelensis*, female, 515 mm TL (WWL 1336); Finca El Acuario, Vereda Las Mercedes, Meta, Colombia, 450 m elevation.



Fig. 489. *Lampropeltis mexicana greeri*; vicinity of Coyotes, Durango, Mexico. Photo by Ed Cassano, courtesy of Louis Porras.



Fig. 490. *Lampropeltis triangulum andesiana*, male, 1,448 mm TL; El Alto del Oso, vicinity of El Parque Nacional Natural La Cueva de los Guácharos, Huila, Colombia, 2,000 m elevation. Photo by John H. Tashjian at the Dallas Zoo.



Fig. 491. *Lampropeltis triangulum arcifera*, male, 890 mm TL (UTA R-12345); 2.4 km northwest of Tapalpa, Jalisco, Mexico, approximately 2,134 m elevation.