

HETERODON PLATIRHINOS (Eastern Hog-nosed Snake). USA: WISCONSIN: PEPIN Co.: specific locality information has been withheld due to the sensitive nature of the site. 27 August 2020. J. Paul White and Heather Kaarakka. Verified by Christopher E. Smith. Milwaukee Public Museum (MPM VZP976a-d; photo vouchers). One juvenile *Heterodon platirhinos* was found within a sand prairie river terrace adjacent to an upland short-grass prairie, areas of open wet meadow and floodplain forest. New county record that fills a gap in the species' documented range in western Wisconsin (Casper 1996. Geographic Distributions of the Amphibians and Reptiles of Wisconsin. Milwaukee Public Museum, Milwaukee, Wisconsin. 87 pp.). The nearest record is ca. 55 km to the southeast in Buffalo County (Becker et al. 2009. Herpetol. Rev. 40:456.).

J. PAUL WHITE (e-mail: john.white@wisconsin.gov), **HEATHER M. KAARAKKA** (e-mail: heather.kaarakka@wisconsin.gov), and **RICHARD A. STAFFEN**, Wisconsin Department of Natural Resources, Bureau of Natural Heritage Conservation, 101 S. Webster St., P.O. Box 7921, Madison, Wisconsin 53707, USA (email: richard.staffen@wisconsin.gov); **JOE E. KRUMRIE** (e-mail: joe.krumrie@wisconsin.gov), **MORGAN G. GEARY** (e-mail: morgan.geary@wisconsin.gov), and **MATT HEREEMA**, Wisconsin Department of Natural Resources, Bureau of Natural Heritage Conservation, 407 S. Second St., Alma, Wisconsin 54610 (email: heeremami30@uw.edu); **ANDREW F. BADJE**, Wisconsin Department of Natural Resources, Bureau of Natural Heritage Conservation, 101 Eagle Dr., Merrill, Wisconsin 54452, USA (e-mail: andrew.badje@wisconsin.gov).

LAMPROPELTIS GREERI (Greer's Kingsnake). MEXICO: ZACATECAS: MUNICIPALITY OF FLORENCIA DE BENITO JUÁREZ: 4.0 km W of Florencia de Benito Juárez (21.49000°N, 103.58800°W; WGS 84), 2197 m elev. 30 May 2020. M. A. Domínguez-de la Riva and B. O. Jiménez-Godoy. Verified by Bradford Hollingsworth. San Diego Society of Natural History, San Diego Natural History Museum (SDSNH Herp PC 05432; photo voucher). The snake was a juvenile found under a rock in a natural grassland with patches of oak forest. First municipality record, bringing the number of municipalities in Zacatecas with records for this species to five (Hansen and Alamillo 2018. Herpetol. Rev. 49:77). The closest reported locality is 21 airline km to the northeast at La Cienega, Municipality of Atolinga, Zacatecas (Ahumada-Carrillo et al. 2011. Herpetol. Rev. 42:397–398).

RUBÉN ALONSO CARBAJAL-MÁRQUEZ, Conservación de la Biodiversidad de México, A.C., Andador Torre de Marfil No. 100, 20229, Aguascalientes, Aguascalientes, Mexico (e-mail: redman031@hotmail.com); **MARCO ANTONIO DOMÍNGUEZ-DE LA RIVA**, Dirección de Regulación Ambiental, Dirección General de Gestión Ambiental, boulevard Torres Landa 1701, 37526, León, Guanajuato, Mexico (e-mail: biol.marcoantonio@gmail.com); **BRYAN OCTAVIO JIMÉNEZ-GODOY**, Universidad Autónoma de Aguascalientes, Centro de Ciencias Básicas, Departamento de Biología, 20131, Aguascalientes, Aguascalientes, Mexico; **GUSTAVO ERNESTO QUINTERO-DÍAZ**, Universidad Autónoma de Aguascalientes, Centro de Ciencias Básicas, Departamento de Biología, 20131, Aguascalientes, Aguascalientes, Mexico (e-mail: gequintmxags@hotmail.com); **JORGE ALBERTO BAÑUELOS-ALAMILLO**, Unidad Académica de Ciencias Biológicas, Universidad Autónoma de Zacatecas, Edificio de Biología Campus II, Ave. Preparatoria S/N, Col. Agronómica, 98066, Zacatecas, Zacatecas, Mexico; Laboratorio de Usos Múltiples, Centro de Bachillerato Tecnológico Agropecuario No. 167, Carretera Agua Fría km 0.5, 99200, Valparaíso, Zacatecas, Mexico (e-mail: j.alberto.ba@gmail.com).

LAMPROPELTIS TRIANGULUM (Milksnake). USA: ILLINOIS: STEPHENSON Co.: east shoulder of Five Corners Rd (42.42291°N,

89.84783°W; WGS 84). 22 August 2020. T. G. Anton. Verified by C. A. Phillips. Illinois Natural History Survey (INHS 44557; photo voucher). Single juvenile found under roadside bark debris. New county record (Smith 1961. Illinois Nat. Hist. Surv. Bull. 28:1–298; Phillips et al. 1999. Field Guide to Amphibians and Reptiles of Illinois. Illinois Natural History Survey Manual 8, Champaign, Illinois. 282 pp.). This record fills an east–west gap between Jo Daviess and Winnebago counties. The nearest record is from 5.0 mi SW of Warren, Jo Daviess County, ca. 10.8 mi (17.5 km) to the northeast (INHS 23450). Specimen observed under an Illinois Department of Natural Resources Research Permit (HSCP 19-44) issued to TGA.

THOMAS G. ANTON, Division of Amphibians and Reptiles, The Field Museum, Roosevelt Road at Lakeshore Drive, Chicago, Illinois 60614, USA.

LAMPROPELTIS TRIANGULUM (Eastern Milksnake). USA: MISSOURI: ADAIR Co.: east side of Spring Lake, 0.6 km W of Hwy N at Whispering Pine Trail (40.10168°N, 92.66598°W; WGS 84). 14 June 2018. Chad E. Montgomery. Verified by Richard Daniel. Dean E. Metter Memorial Collection, University of Missouri, Columbia (UMC 3475P; photo voucher). Sub-adult male (55.0 cm SVL, 10.2 cm tail length, 98.1 g). New county record in Missouri, ca. 64 km SW of the closest record in Scotland County, Missouri (UMC 2055P; Daniel and Edmond 2018. Atlas of Missouri Reptiles and Amphibians for 2017. <http://atlas.moherp.org/pubs/atlas17.pdf>; 15 June 2018).

CALVIN H. SCHAEFER (e-mail: chs8581@truman.edu) and **CHAD E. MONTGOMERY**, Biology Department, Truman State University, Kirksville, Missouri, 63501, USA (e-mail: chadmont@truman.edu).

LATICAUDA COLUBRINA (Yellow-lipped Sea Krait). PHILIPPINES: BIRI ISLAND: NORTHERN SAMAR PROVINCE: MUNICIPALITY OF BIRI: Biri-Larosa Protected Landscape and Seascape (12.68176°N, 124.36037°E; WGS 84). 13 July 2015. Rowell P. Abogado. Verified by Wolfgang Wüster. Lee Kong Chian Natural History Museum, National University of Singapore (ZRC [IMG] 2.546; photo voucher). Individual found swimming through seagrass bed ca. 50 m from shore of Barangay Poblacion, ca. 2 m below sea level. This species has been documented on many islands and rocky, coastal shores of the Philippines, including southern Luzon, Hundred Islands, Palawan, islands in Central Visayas, Gigantes Islands, Zamboanga peninsula, Bongo Island in Tawi-tawi, Sulu Islands (Heatwole et al. 2005. Herpetol. Monogr. 19:1–136), and Maestre de Campo Island in Romblon (Siler et al. 2012. Check List 8:443–462). The nearest known, but old, record is ca. 176 km to the south-southeast in the coastal village of Nipa-nipa (now defunct and integrated into Municipality of Marabut, Province of Samar), southwest Samar (Peters 1862. Monatsb. K. Preuss. Akad. Wiss. Berlin 1861:683–691). First record for island, municipality, and province, shortening the distribution gap between provinces of Albay and Samar in the eastern Philippines.

ROWELL P. ABOGADO, Barangay Calo, Bay, Laguna 4033, Philippines (e-mail: rpabogado@up.edu.ph); **CYRUS JOB P. DELA CRUZ**, Department of Environment and Natural Resources – Regional Office V, Rawis, Legazpi City, Albay 4500, Philippines (e-mail: cyrusjob.delacruz@icol-u.edu.ph).

LATICAUDA LATICAUDATA (Blue-lipped Sea Krait). PHILIPPINES: BIRI ISLAND: NORTHERN SAMAR PROVINCE: MUNICIPALITY OF BIRI: Biri-Larosa Protected Landscape and Seascape (12.68616°N, 124.38182°E; WGS 84), 0 m elev. 12 July 2015. Rowell P. Abogado. Verified by Wolfgang Wüster. Lee Kong Chian Natural