LAMPROPELTIS ALTERNA (Gray-banded Kingsnake). MEXICO: DURANGO: Municipio de Santiago Papasquiaro: East slope of the Sierra Madre Occidental (25.087°N, 105.542°W, WGS 84), 2218 m elev. 31 July 2007. J. Reyes Velasco and M. J. Ingrasci. Verified by J. A. Campbell. University of Texas Arlington Digital Collection (UTADC 1096). New municipality

record, third record for Durango, and first record for Sierra Madre Occidental, extending the known range ca. 100 km W of the closest other known Durango record, 5 mi. S of San Antonio near El Banco; 26 mi. N San Juan del Río, 4700 ft. elev. (Tanzer 1970. Herpetologica 26:419–428 [as *L. mexicana*]). Garstka (1982. Breviora 466:1–35 [as *L. alterna*]) reported the same specimen using the 26 mi. N San Juan Del Río locality (TCWC 36892) and Gadsden et al. (2006. Bull. Chicago Herpetol. Soc. 41:2–9) reported another taken in the Sierra el Sarnoso, Municipio Lerdo (between 1100 m and 1350 m elev.) in the northeastern portion of the state. The other two records are from Chihuahuan Desert scrub vegetation, whereas the snake reported herein was found in Madrean woodland vegetation at a higher elevation.

Submitted by MATTHEW J. INGRASCI (e-mail: mingrasc@hotmail.com); KIRK SETSER, Departamento de Biología Animal, Universidad de Granada, 18071, Granada, Spain (e-mail: kwsetser@gmail.com); and JACOBO REYES VELASCO, Calle del Parque # 45, Colonia Chapalita, Zapopan, Jalisco, Mexico 45042 (e-mail: jackobz@gmail.com).