

LAMPROPELTIS MEXICANA (San Luis Potosi Kingsnake). MÉXICO: NUEVO LEÓN: 8.6 km E Iturbide on Rt. 58, Municipio Iturbide ($24^{\circ}44'45.5''N$, $99^{\circ}50'61.6''W$; 1036 m). 24 July 2000. Gerard T. Salmon, Daniel Vermilya, and Bartholomew B. Bruno. Universidad Autónoma de Nuevo León Herpetological Collection (UANL 5773). Verified by Hobart M. Smith. DOR adult male. Easternmost locality for the species in the state of Nuevo León. Extends reported range ca. 19 air km to the east of TU 16483 (Garstka 1982, Breviora 466:1–35) and to E slope of the Sierra Madre Oriental. Lowest elevation record for the species (Gehlbach 1967, Cat. Am. Amphib. Rept. 55).

Submitted by **GERARD T. SALMON**, 30 Appletree Drive, Rhinebeck, New York 12572, USA, **ROBERT W. BRYSON, JR.**, Department of Biology, Sul Ross State University, Alpine, Texas 79832, USA, and **DAVID LAZCANO**, Laboratorio de Herpetología, Universidad Autónoma de Nuevo León, Apartado Postal 513, San Nicolas de los Garza, Nuevo León, Caixa Postal 66450, México.

LAMPROPELTIS MEXICANA (San Luis Potosi Kingsnake). MÉXICO: SINALOA: 4 km W El Palmito on Rt. 40, Municipio Concordia ($23^{\circ}33'14.2''N$, $105^{\circ}50'47.2''W$, 2000 m). 30 June 2000. Robert Bryson, Deron Hartman, and Javier Banda. Universidad Autónoma de Nuevo León Herpetological Collection (UANL 5684). Verified by Robert G. Webb. DOR juvenile. First state record and westernmost locality for species. Extends reported range ca. 95 air km to the west (Webb 1984, In Seigel et al. [eds.], Univ. Kansas Mus. Nat. Hist. Spec. Publ. 10:1–278) and ca. 10 air km from a previously unreported specimen of *L. mexicana* (FWMSH 6716) from Durango.

Submitted by **ROBERT W. BRYSON, JR.**, Department of Biology, Sul Ross State University, Alpine, Texas 79832, USA, **GERARD T. SALMON**, 30 Appletree Drive, Rhinebeck, New York 12572, USA, and **DAVID LAZCANO**, Laboratorio de